

Fig.1




Fig. 2/




Fig. 2//




Fig. 3/




Fig. 3//




Fig.4/




Fig. 4 //




Fig.5




Fig. 6/




Fig. 6//




Fig. 7




Fig. 8A'




Fig. 8A //




Fig. 8B/




Fig. 8B //




Fig. 9A'




Fig. 9A //




Fig. 9A ///




Fig. 9B/




Fig. 9B//




Fig. 9B///




Fig. 9C/




Fig. 9C //




Fig. 9C///






Fig. 9D/








Fig. 9E/




Fig. 9E //




Fig.9E///




Fig. 10A'




Fig. 10A //




Fig. 10B/





Fig. 10B //






Fig. 10C




Fig. 10D/






Fig. 10D








Fig. 10F/




Fig. 10F //




Fig. 11: Human Cystatin C immunodetection in CSF and plasma samples





Fig.12: Bovine haemoglobin immunodetection in plasma samples



10-20% Tricine Gels Gradipore

Ac I: Affinity pure goat anti Hb human, Biomeda 1/100 (10 ug/ml)

Ac II: Rabbit Ig G anti goat HRP, Dako, 1/2000

bovine haemoglobin: 100 ng of purified proteins from bovine erythrocytes

Human haemoglobin: 100 ng of semi-pure oxyhaemoglobin from human erythrocytes

Fig.13: Analytical silver nitrate stained 2D gel of BSE - plasma sample




Fig. 14: SELDI profiling of human and bovine Haemoglobin
on NP20 (native protein 66 kda)




Fig. 15A: Comparison of bovine Hb Seldi profile and BSE + plasma samples




Fig.15B: Comparison of bovine Hb Seldi profile and
BSE + plasma samples between 7 and 50 kda




Fig.15C: Comparison of bovine Hb Seldi profile and BSE + plasma sample < 7 kda

